

# BABAR status and prospects for $CP$ asymmetry measurements: $\sin(2\beta + \gamma)$

S. Ganzhur\*

DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France

The recent experimental results on  $CP$  violation related to the angles of the Cabibbo-Kobayashi-Maskawa (CKM) unitarity triangle  $2\beta + \gamma$  are summarized in these proceedings. These results are obtained with approximately 232 million  $\Upsilon(4S) \rightarrow B\bar{B}$  events collected with the BABAR detector at the PEP-II asymmetric-energy  $B$ -factory at SLAC. Using the measurements on time-dependent  $CP$  asymmetries in  $B^0 \rightarrow D^{(*)\mp}\pi^\pm$  and  $B^0 \rightarrow D^\mp\rho^\pm$  decays and theoretical assumptions, one finds  $|\sin(2\beta + \gamma)| > 0.64$  (0.40) at 68% (90%) confidence level. The perspectives of  $\sin(2\beta + \gamma)$  measurement with  $\bar{B}^0 \rightarrow D^{(*)0}\bar{K}^{(*)0}$  and  $B^0 \rightarrow D^{(*)\mp}a_{0(2)}^\pm$  decay channels are also discussed.

## Introductoin

A crucial part of the  $CP$  violation program in  $B$ -factories is the measurement of the angle  $\gamma(\phi_3) = \arg(-V_{ud}V_{ub}^*/V_{cd}V_{cb}^*)$  of the unitary triangle related to the CKM matrix [1]. Decays of  $B_d$  mesons that allows one to constraint the CKM angle  $2\beta + \gamma$ , have either small  $CP$  asymmetry ( $B \rightarrow D^{(*)}\pi$ ) or small branching fractions ( $B \rightarrow D^{(*)}K^{(*)}$ ). This makes the  $CP$  violation effect hard to measure. Furthermore, due to presence of two hadronic parameters in the observables ( $r$  and  $\delta$ , the amplitude ratio and the strong phase difference between two amplitudes) it is difficult to cleanly extract the weak phase information, although approaches based on  $SU(3)$  symmetry exists.

## I. THE BABAR DETECTOR AND DATASET

The data used in the presented analyzes were recorded with the BABAR detector at the PEP-II asymmetric-energy storage rings, and consist of  $211 \text{ fb}^{-1}$  collected on the  $\Upsilon(4S)$  resonance (on-resonance sample), and  $21 \text{ fb}^{-1}$  collected at an  $e^+e^-$  center-of-mass (CM) energy approximately 40 MeV below the resonance peak (off-resonance sample). This corresponds to approximately 232 million  $\Upsilon(4S) \rightarrow B\bar{B}$  recorded events.

The BABAR detector is described in detail in Ref. [2].

## II. $CP$ ASYMMETRY IN $B^0 \rightarrow D^{(*)\mp}\pi^\pm/\rho^\pm$ DECAYS

The decay modes  $B^0 \rightarrow D^{(*)\mp}\pi^\pm$  have been proposed to measure  $\sin(2\beta + \gamma)$  [3]. In the Standard Model the decays  $B^0 \rightarrow D^{(*)+}\pi^-$  and  $\bar{B}^0 \rightarrow D^{(*)-}\pi^+$  proceed through the  $\bar{b} \rightarrow \bar{u}cd$  and  $b \rightarrow c$  amplitudes  $A_u$  and  $A_c$ , respectively. The relative weak phase between these two amplitudes is  $\gamma$ . When combined with  $B^0\bar{B}^0$  mixing,

this yields a weak phase difference of  $2\beta + \gamma$  between the interfering amplitudes.

The decay rate distribution for  $B \rightarrow D^{(*)\pm}\pi^\mp$  is

$$P_\eta^\pm(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} \times [1 \mp S^\zeta \sin(\Delta m \Delta t) \mp \eta C \cos(\Delta m \Delta t)], \quad (1)$$

where  $\tau$  is the  $B^0$  lifetime averaged over the two mass eigenstates,  $\Delta m$  is the  $B^0 - \bar{B}^0$  mixing frequency, and  $\Delta t$  is the difference between the time of the  $B \rightarrow D^{(*)\pm}\pi^\mp$  ( $B_{\text{rec}}$ ) decay and the decay of the other  $B$  ( $B_{\text{tag}}$ ) in the event. The upper (lower) sign in Eq. 1 indicates the flavor of the  $B_{\text{tag}}$  as a  $B^0$  ( $\bar{B}^0$ ), while  $\eta = +1$  ( $-1$ ) and  $\zeta = +$  ( $-$ ) for the  $B_{\text{rec}}$  final state  $D^{(*)-}\pi^+$  ( $D^{(*)+}\pi^-$ ). The parameters  $C$  and  $S^\pm$  are given by

$$C \equiv \frac{1 - r^2}{1 + r^2}, \quad S^\pm \equiv \frac{2r}{1 + r^2} \sin(2\beta + \gamma \pm \delta). \quad (2)$$

Here  $\delta$  is the strong phase difference between  $A_u$  and  $A_c$  and  $r \equiv |A_u/A_c|$ . Since  $A_u$  is doubly CKM-suppressed with respect to  $A_c$ , one expects  $r$  to be small of order 2%. Due to the small value of  $r$ , large data samples are required for a statistically significant measurement of  $S$ .

Since the expected  $CP$  asymmetry in the selected  $B$  decays is small, this measurement is sensitive to the interference between the  $b \rightarrow u$  and  $b \rightarrow c$  amplitudes in the decay of  $B_{\text{tag}}$ . To account for this “tag-side interference”, we use a parametrization which is described in Ref. [4]. The  $S^\pm$  coefficient are replaced with three others

$$\begin{aligned} a &= 2r \sin(2\beta + \gamma) \cos \delta \\ b &= 2r' \sin(2\beta + \gamma) \cos \delta' \\ c &= 2 \cos(2\beta + \gamma) (r \sin \delta - r' \sin \delta') \end{aligned} \quad (3)$$

For each tagging category, independent of the decay mode  $\{D\pi, D^*\pi, D\rho\}$ , the tag-side interference is parametrized in terms of the effective parameters  $r'$  and  $\delta'$ . One notes,  $r' = 0$  for the lepton tagging category.

Two different analysis techniques, full reconstruction [5] and partial reconstruction [6] were used for the  $\sin(2\beta + \gamma)$  measurement with  $B^0 \rightarrow D^{(*)\mp}\pi^\pm$ .

The full reconstruction technique is used to measure the  $CP$  asymmetry in  $B^0 \rightarrow D^{(*)\mp}\pi^\pm$  and  $B^0 \rightarrow D^{*\mp}\rho^\pm$

\*Electronic address: ganzhur@cea.fr

decays [7]. From a time-dependent maximum likelihood fit the following parameters related to the  $CP$  violation angle  $2\beta + \gamma$  are obtained:

$$\begin{aligned} a^{D\pi} &= -0.010 \pm 0.023 \pm 0.007 \\ c_{\text{lep}}^{D\pi} &= -0.033 \pm 0.042 \pm 0.012 \\ a^{D^*\pi} &= -0.040 \pm 0.023 \pm 0.010 \\ c_{\text{lep}}^{D^*\pi} &= 0.049 \pm 0.042 \pm 0.015 \\ a^{D\rho} &= -0.024 \pm 0.031 \pm 0.009 \\ c_{\text{lep}}^{D\rho} &= -0.098 \pm 0.055 \pm 0.018 \end{aligned} \quad (4)$$

where the first error is statistical and the second is systematic. The systematic error for  $B^0 \rightarrow D^{*\mp}\pi^\pm$  includes the maximum bias of asymmetry parameters due to possible dependence of  $r$  on the  $\pi\pi^0$  invariant mass. For the measurement of  $2r \cos(2\beta + \gamma) \sin \delta$  parameter only the lepton-tagged events are used due to a presence of tag-side  $CP$  violation effect [4].




FIG. 1: Raw asymmetry for (a) lepton-tagged and (b) kaon-tagged events of  $B^0 \rightarrow D^{*\mp}\pi^\pm$  decay mode using the method of the partial reconstruction. The curves represent the projections of the PDF for the raw asymmetry.

In the partial reconstruction of a  $B^0 \rightarrow D^{*\mp}\pi^\pm$  candidate, only the hard (high-momentum) pion track  $\pi_h$  from the  $B$  decay and the soft (low-momentum) pion track  $\pi_s$  from the decay  $D^{*-} \rightarrow \bar{D}^0\pi_s^-$  are used. Applying kinematic constraints consistent with the signal decay mode, the four-momentum of the non-reconstructed, “missing”  $D$  is calculated. Signal events are peaked in the  $m_{\text{miss}}$  distribution at the nominal  $D^0$  mass. This method eliminates the efficiency loss associated with the neutral  $D$  meson reconstruction. The  $CP$  asymmetry independent on the assumption on  $r^{D^*\pi}(r^*)$  measured with this technique is [8]

$$\begin{aligned} a^{D^*\pi} &= -0.034 \pm 0.014 \pm 0.009 \\ c_{\text{lep}}^{D^*\pi} &= -0.019 \pm 0.022 \pm 0.013 \end{aligned} \quad (5)$$

where the first error is statistical and the second is systematic. This measurement deviates from zero by 2.0 standard deviations. Figure 1 shows the raw, time-dependent  $CP$  asymmetry

$$A(\Delta t) = \frac{N_{B^0}(\Delta t) - N_{\bar{B}^0}(\Delta t)}{N_{B^0}(\Delta t) + N_{\bar{B}^0}(\Delta t)} \quad (6)$$

In the absence of background and with high statistics, perfect tagging, and perfect  $\Delta t$  measurement,

$A(\Delta t)$  would be a sinusoidal oscillation with amplitude  $2r \sin(2\beta + \gamma) \cos \delta$ .

Two methods for interpreting these results in terms of constraints on  $|\sin(2\beta + \gamma)|$  are used. Both methods involve minimizing a  $\chi^2$  function that is symmetric under the exchange  $\sin(2\beta + \gamma) \rightarrow -\sin(2\beta + \gamma)$ , and applying the method of Ref. [9]. In the first interpretation method, no assumption regarding the value of  $r^*$  is made. The resulting 95% lower limit for the mode  $B^0 \rightarrow D^{*\mp}\pi^\pm$  is shown as a function of  $r^*$  in Figure 2. The second interpretation assumes that  $r^{(*)}$  can be estimated from the Cabibbo angle, the ratio of branching fractions  $\mathcal{B}(B^0 \rightarrow D^{(*)+}_s \pi^-)/\mathcal{B}(B^0 \rightarrow D^{(*)-} \pi^+)$ , and the ratio of decay constants  $f_{D^*}/f_{D_s^*}$ . The confidence level as a function of  $|\sin(2\beta + \gamma)|$  is shown in Figure 3. This method yields the lower limits  $|\sin(2\beta + \gamma)| > 0.64$  (0.40) at 68% (90%) C.L.




FIG. 2: Lower limit on  $|\sin(2\beta + \gamma)|$  at 90% CL as a function of  $r^{(*)}$ , for  $r^{(*)} > 0.001$ .




FIG. 3: The shaded region denotes the allowed range of  $|\sin(2\beta + \gamma)|$  for each confidence level. The horizontal lines show, from top to bottom, the 68% and 90% CL.

### III. $\bar{B}^0 \rightarrow D^{(*)0} \bar{K}^{*0}$ DECAYS

The decay modes  $\bar{B}^0 \rightarrow D^{(*)0} \bar{K}^0$  have been proposed for determination of  $\sin(2\beta + \gamma)$  from measurement of time-dependent  $CP$  asymmetries [10]. In the Standard Model the decays of  $B^0$  and  $\bar{B}^0$  mesons into final state  $D^{(*)0} K_s^0$  proceed through the  $b \rightarrow c$  and  $\bar{b} \rightarrow \bar{u}$  amplitudes, respectively. Due to relatively large  $CP$  asymmetry ( $r_B \equiv |A(\bar{B}^0 \rightarrow \bar{D}^{(*)0} \bar{K}^0)|/|A(\bar{B}^0 \rightarrow D^{(*)0} \bar{K}^0)| \simeq 0.4$ ) these decay channels look very attractive for such a measurement. Since the parameter  $r_B$  can be measured with sufficient data sample by fitting the  $C$  coefficient in time distributions, the measured asymmetry can be interpreted in terms of  $\sin(2\beta + \gamma)$  without additional assumptions. However, the branching fractions of such decays are relatively small ( $\sim 5 \cdot 10^{-5}$ ). That is way the large data sample is still required.

From the measured signal yields [11], we find

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^0 \bar{K}^0) &= (5.3 \pm 0.7 \pm 0.3) \times 10^{-5} \\ \mathcal{B}(\bar{B}^0 \rightarrow D^{*0} \bar{K}^0) &= (3.6 \pm 1.2 \pm 0.3) \times 10^{-5} \\ \mathcal{B}(\bar{B}^0 \rightarrow D^0 \bar{K}^{*0}) &= (4.0 \pm 0.7 \pm 0.3) \times 10^{-5} \\ \mathcal{B}(\bar{B}^0 \rightarrow \bar{D}^0 \bar{K}^{*0}) &< 1.1 \times 10^{-5} \text{ at 90\% C.L.} \end{aligned} \quad (7)$$

where the uncertainties are statistical and systematic, respectively. Figure 4 shows the  $\Delta E$  distributions of candidates with  $|m_{\text{ES}} - 5280| < 8 \text{ MeV}/c^2$  for the sums of the reconstructed  $D^0$  decay modes.




FIG. 4: Distribution of  $\Delta E$  for a)  $\bar{B}^0 \rightarrow D^0 \bar{K}^0$ , b)  $\bar{B}^0 \rightarrow D^{*0} \bar{K}^0$ , c)  $\bar{B}^0 \rightarrow D^0 \bar{K}^{*0}$ , and d)  $\bar{B}^0 \rightarrow \bar{D}^0 \bar{K}^{*0}$  candidates with  $|m_{\text{ES}} - 5280 \text{ MeV}/c^2| < 8 \text{ MeV}/c^2$ . The points are the data, the solid curve is the projection of the likelihood fit, and the dashed curve represents the background component.

The  $B$  decay dynamics can modify the expectation for the ratio  $r_B$ . The magnitude of this ratio can be probed by measuring the rate for the decays  $\bar{B}^0 \rightarrow D^{(*)0} \bar{K}^{*0}$  and  $\bar{B}^0 \rightarrow \bar{D}^{(*)0} \bar{K}^{*0}$  using the self-tagging decay  $\bar{K}^{*0} \rightarrow K^- \pi^+$ . The  $\bar{B}^0 \rightarrow D^0 \bar{K}^{*0}$  and  $\bar{B}^0 \rightarrow \bar{D}^0 \bar{K}^{*0}$  decays are distinguished by the correlation between the charges of

the kaons produced in the decays of the neutral  $D$  and the  $\bar{K}^{*0}$ . This charge correlation in the final state is diluted by the presence of the doubly-Cabibbo-suppressed decays  $D^0 \rightarrow K^+ \pi^-$ ,  $K^+ \pi^- \pi^0$ , and  $K^+ \pi^- \pi^+ \pi^-$ . The ratio  $r_B$  is related to the experimental observable  $\mathcal{R}$  defined for the  $D^0 \rightarrow K^+ \pi^-$  decay as

$$\begin{aligned} \mathcal{R} &= \frac{\Gamma(\bar{B}^0 \rightarrow (K^+ \pi^-)_D \bar{K}^{*0})}{\Gamma(\bar{B}^0 \rightarrow (K^- \pi^+)_D \bar{K}^{*0})} \\ &= r^{(*)2}_B + r_D^2 + 2r^{(*)}_B r_D \cos(\gamma + \delta), \end{aligned} \quad (8)$$

where

$$r_D = \frac{|\mathcal{A}(D^0 \rightarrow K^+ \pi^-)|}{|\mathcal{A}(D^0 \rightarrow K^- \pi^+)|}, \quad (9)$$

$$\delta = \delta_B + \delta_D, \quad (10)$$

and  $\delta_B$  and  $\delta_D$  are strong phase differences between the two  $B$  and  $D$  decay amplitudes, respectively. From the measured  $B$  branching fractions (Eq. 7), values of  $r_D$  [12] and Eq. 8, one obtains  $r < 0.40$  at the 90% C.L. To conclude, the present signal yields combined with this limit on  $r$  suggest that a substantially larger data sample is needed for a competitive time-dependent measurement of  $\sin(2\beta + \gamma)$  in  $\bar{B}^0 \rightarrow D^{(*)0} \bar{K}^0$  decays.

### IV. $B^0 \rightarrow D^{(*)\mp} a_{0(2)}^\pm$ DECAYS

Recently it was proposed to consider the  $B^0 \rightarrow D^{(*)\mp} a_{0(2)}^\pm$  decays for measurement of  $\sin(2\beta + \gamma)$  [13]. The decay amplitudes of  $B$  mesons to light scalar or tensor mesons such as  $a_0^+$  or  $a_2^+$ , emitted from a weak current, are significantly suppressed due to the small decay constants  $f_{a_{0(2)}}$ . Thus, the absolute value of the CKM-suppressed and favored amplitudes become comparable. As a result, the  $CP$  asymmetry in such decays is expected to be large. However, the theoretical predictions of the branching fractions for  $B^0 \rightarrow D^{(*)\mp} a_{0(2)}^\pm$  is expected of the order of  $(1 \div 4) \cdot 10^{-6}$  [14]. The main uncertainty in the branching fractions of these decay modes is due to unknown  $B \rightarrow a_{0(2)} X$  transition form factors. One way to verify the expectations and test a validity of the factorization approach is to measure the branching fractions for the more abundant decay modes  $B^0 \rightarrow D_s^{(*)+} a_{0(2)}^-$ .

Using a sample of about 230 million  $\Upsilon(4S) \rightarrow B\bar{B}$  no evidence for these decays were observed [15]. This allowed one to set upper limits at 90% C.L. on the branching fractions to be

$$\begin{aligned} \mathcal{B}(B^0 \rightarrow D_s^+ a_0^-) &< 1.9 \cdot 10^{-5} \\ \mathcal{B}(B^0 \rightarrow D_s^+ a_2^-) &< 1.9 \cdot 10^{-4} \\ \mathcal{B}(B^0 \rightarrow D_s^{*+} a_0^-) &< 3.6 \cdot 10^{-5} \\ \mathcal{B}(B^0 \rightarrow D_s^{*+} a_2^-) &< 2.0 \cdot 10^{-4} \end{aligned} \quad (11)$$

Figure 5 shows the  $m_{\text{ES}}$  distributions for the reconstructed candidates  $B^0 \rightarrow D_s^+ a_0^-$ ,  $B^0 \rightarrow D_s^+ a_2^-$ ,  $B^0 \rightarrow$




FIG. 5: Distributions of  $m_{ES}$  for a)  $B^0 \rightarrow D_s^+ a_0^-$ , b)  $B^0 \rightarrow D_s^+ a_2^-$ , c)  $B^0 \rightarrow D_s^{*+} a_0^-$ , d)  $B^0 \rightarrow D_s^{*+} a_2^-$  candidates overlaid with the projection of the maximum likelihood fit. Contributions from the three  $D_s^+$  decay modes are shown with different hatching styles:  $\phi\pi^+$  is cross hatched,  $\bar{K}^{*0}K^+$  is hatched, and  $K_s^0 K^+$  is white.

$D_s^{*+} a_0^-$  and  $B^0 \rightarrow D_s^{*+} a_2^-$ . For each  $B$  decay mode, an unbinned maximum-likelihood fit is performed using the candidates from the three  $D_s^+$  decay modes.

The upper limit value for  $B^0 \rightarrow D_s^{*+} a_0^-$  is lower than the theoretical expectation, which might indicate the need to revisit the  $B \rightarrow a_0 X$  transition form factor estimate. It might also imply the limited applicability of the factorization approach for this decay mode. The measured upper limits suggest that the branching ratios of  $B^0 \rightarrow D^{(*)+} a_{0(2)}^-$  are too small for  $CP$ -asymmetry mea-

surements given the present statistics of the  $B$ -factories. The measurement of  $\sin(2\beta + \gamma)$  in  $B^0 \rightarrow D^{(*)+} a_{0(2)}^-$  decays is an interesting program for the future experiments such as Super $B$ -factories.

## Conclusion

The substantial constraint on the CKM angles  $2\beta + \gamma$  comes from the measurements of time-dependent  $CP$  asymmetry in the  $B^0 \rightarrow D^{(*)\mp} \pi^\pm$  and  $B^0 \rightarrow D^\mp \rho^\pm$  decays. The *BABAR* experiment has used two techniques such as full and partial reconstruction to increase the signal yields in the  $D^{*\mp} \pi^\pm$  channel. The combined *BABAR* and *BELLE* results [16] for  $CP$  violation in the most precisely measured decay channel  $D^{*\mp} \pi^\pm$  is

$$a^{D^* \pi} = 2r^* \sin(2\beta + \gamma) \cos \delta = -0.037 \pm 0.011 \quad (12)$$

This measurement performed at the level of one per cent deviates from zero by 3.4 standard deviations. Future updates are therefore of a great interest. We interpret the *BABAR* result in terms of  $\sin(2\beta + \gamma)$  and find  $|\sin(2\beta + \gamma)| > 0.64$  (0.40) at 68% (90%) C.L. using a frequentist method.

The *BABAR* experiment has measured the branching fractions of  $\bar{B}^0 \rightarrow D^{(*)0} \bar{K}^{(*)0}$  and set up the limit on  $B^0 \rightarrow D^{(*)\mp} a_{0(2)}^\pm$  decays. The present signal yields and established limits suggest that a substantially larger data sample is needed for a competitive time-dependent measurement of  $\sin(2\beta + \gamma)$  with these decay channels.

- 
- [1] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
  - [2] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instrum. Meth. A **479**, 1 (2002).
  - [3] R.G. Sachs, Enrico Fermi Institute Report, EFI-85-22 (1985) (unpublished); I. Dunietz and R.G. Sachs, Phys. Rev. **D37**, 3186 (1988) [E: Phys. Rev. **D39**, 3515 (1989)]; I. Dunietz, Phys. Lett. **B427**, 179 (1998); P.F. Harrison and H.R. Quinn, ed., “The *BABAR* Physics Book”, SLAC-R-504 (1998), Chap. 7.6.
  - [4] O. Long, M. Baak, R.N. Cahn, and D. Kirkby, Phys. Rev. D **68**, 034010 (2003).
  - [5] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **92**, 251801 (2004).
  - [6] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **92**, 251802 (2004).
  - [7] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **73**, 111101 (2006).
  - [8] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **71**, 112003 (2005).
  - [9] G. Feldman and R. Cousins, Phys. Rev. D **57**, 3873 (1998).
  - [10] M. Gronau and D. London, Phys. Lett. B **253**, 483 (1991); D. Atwood, I. Dunietz, and A. Soni, Phys. Rev. Lett. **78**, 3257 (1997); B. Kayser and D. London, Phys. Rev. D **61**, 116013 (2000); A.I.Sandra, hep-ph/0108031.
  - [11] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **74** 031101 (2006)
  - [12] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592** 1 2004 .
  - [13] M. Diehl, G. Hiller, Phys. Lett. B **517**, 125 (2001).
  - [14] M. Diehl, G. Hiller, “New ways to explore factorization in  $B$  decays”, hep-ph/0105194
  - [15] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **73** 071103 (2006)
  - [16] *BELLE* Collaboration, F.J. Ronga *et al.*, Phys. Rev. D **73**, 092003 (2006).